arXiv:hep-ph/0508261v1 25 Aug 2005

A COMPREHENSIVE APPROACH TO STRUCTURE FUNCTIONS [1]

A. DONNACHIE

School of Physics and Astronomy, University of Manchester,
Manchester M13 9PL, England

Abstract

A model is presented based on a dipole picture with a hard and a soft pomeron. It is assumed that large dipoles couple to the soft pomeron and small dipoles couple to the hard pomeron. Most of the parameters of the model are predetermined from proton-proton scattering and the only free parameter is the radius R_c, defining the transition from small to large dipoles. This is fixed by the proton structure function $F_2(x,Q^2)$. The model then successfully predicts $F_2^c(x,Q^2)$, $F_2^L(x,Q^2)$, J/ψ photoproduction, $\gamma^* p \to \gamma p$, $\sigma_{\gamma p}^{\rm Tot}(s)$, $\sigma_{\gamma\gamma}^{\rm Tot}(s)$ and $F_2^\gamma(x,Q^2)$.

1 Objective

Deep inelastic scattering at small x can be described successfully by a two-component model [1] comprising the soft non-perturbative pomeron of hadronic interactions, intercept $\sim 0.08-0.1$, and a hard pomeron, intercept ~ 1.4. Although phenomenologically successful, it does not explain, for example, the relative strengths of the hard and soft pomeron in deep inelastic scattering and in J/ψ photoproduction, or why the charm structure function of the proton is dominated by the hard pomeron.

To answer these questions requires a specific model for the diffractive process, consideration of the particle wave functions and simultaneous treatment of several reactions to separate the dynamics of diffraction from wave-function effects. The intent is to obtain a global description of a variety of data in a simple model, not necessarily to obtain a detailed fit, although in practice the model is surprisingly successful despite its simplicity.

2 The model

The model proposed [2] is a simple one, based on a dipole picture with two pomerons in which small dipoles couple to the hard pomeron and large dipoles couple to the soft pomeron. Hadron-hadron, photon-hadron and photon-photon reactions are treated in a uniform approach. This is based on the "Heidelberg Model" [3] of soft diffraction and the proton is considered as a

---

[1] Summary of talk given at the XIth International Conference on Elastic and Diffractive Scattering, Blois 2005



Figure 1: Fit to $\sigma^{\mathrm{Tot}}_{\gamma^* p}(Q^2, W)$ for different values of Q^2.The data are from ZEUS [6] (squares) and H1 [7] (triangles).

quark-diquark system, that is effectively as a dipole. The parameters of the photon and proton wave functions and the dipole-dipole cross section are taken from applications of that model to soft hadronic [4] and photoproduction reactions [5]. These parameters remain unchanged throughout, the criteria for defining small and large dipoles being obtained from the proton structure function. All other processes are controlled by the relevant particle wave functions.

The standard perturbative expressions were used for the photon wave functions with a Q^2-dependent quark mass $m_{eff}(Q^2)$ for photons of low virtuality (including real).

$$\begin{array}{rlll} m_{eff}(Q^2) = & m_f & +m_{0q}(1 - Q^2/Q_0^2) & Q^2 \leq Q_0^2, \\ & m_f & & Q^2 \geq Q_0^2. \end{array} \qquad (1)$$

The masses used were $m_{0q} = 0.19$ GeV for the light quarks, 0.31 GeV for the strange quark and 1.25 GeV for the charm quark.

A Gaussian wave function was assumed for the proton:

$$\psi_p(\vec{R}) = \frac{1}{\sqrt{2\pi}} exp\Big(- \frac{R^2}{4R_p^2}\Big). \tag{2}$$

The radius R_p was chosen to reproduce the logarithmic slope of elastic pp scattering at $\sqrt{s} = 20$ GeV, giving $R_p = 0.75$ fm. The J/ψ wave function is taken as that of a massive vector current with mass m_c and with the radial dependence modelled by a Gaussian, the radius parameter being fixed by the electromagnetic decay width. Full details can be found in [5].

The Heidelberg Model is defined at $W = \sqrt{s} = 20$ GeV, so energy dependence has to be introduced by hand by dividing the amplitude into a soft and a hard part:

$$T_{ab\to cd} = iW^2 \Big(T^{\rm soft}_{ab\to cd} (W/W_0)^{2\epsilon_s} + T^{\rm hard}_{ab\to cd}(W/W_0)^{2\epsilon_h} \Big) \tag{3}$$

with $W_0 = 20$ GeV, $\epsilon_s = 0.08$ and $\epsilon_h = 1.42$.

To be economical with parameters a sharp cut was introduced, assuming that only the soft pomeron couples if both dipoles are larger than a certain value R_c whereas the hard pomeron couples if at least one of the dipoles is smaller than R_c.

3 $\gamma^* p$ reactions

The parameter R_c, the only free parameter in the model, was obtained by fitting the proton structure function, or rather the total $\gamma^* p$ cross section, from $Q^2 \approx 0$ to 150 GeV^2. The result is $R_c = 0.22$ fm and examples of the fit are shown in figure 1.

With R_c fixed, $F_2^c(x, Q^2)$ and $F_L(x, Q^2)$ can be predicted and agreement with the corresponding data is excellent. The model is compared with the F_2^c data in figure 2. In both F_2^c and F_2^L the photon wave function is concentrated at smaller distances than in F_2, so the hard pomeron is already dominant at moderate energies. Although the increase of the hard-pomeron contribution in F_L with increasing Q^2 is not quite as strong as in F_2 the soft-pomeron contribution is even more suppressed so the hard pomeron is dominant sooner in F_L than in F_2. We emphasize that these are purely wave-function effects. The model provides an explanation for the phenomenological result [10] that $F_2^c(x, Q^2)$ is completely dominated by the hard pomeron, as can be seen in figure 2.

The ratio of the soft pomeron to the hard pomeron in J/ψ photoproduction is much larger than in F_2^c at comparable energies. This is again a wave-function effect. The virtual $c\bar{c}$ pair in the photon wave function has an extension $\approx 1/m_c$. The J/ψ has a much larger radius, so the overlap of the charm part of the photon wave function with the J/ψ wave function obtains a larger contribution from $R > R_c$ than does the square of the charm part of the photon wave function. The model describes the J/ψ total cross section well [2], either assuming a constant logarithmic slope of $b = 6$ GeV^{-2} or with a slope varying with energy as predicted by Regge theory. For the latter the slope of the soft-pomeron trajectory was taken as the standard $\alpha_{P_s} = 0.25$ GeV^{-2} and that of the hard-pomeron trajectory as $\alpha_{P_h} = 0.1$ GeV^{-2} with $b = 6$ GeV^{-2} at $\sqrt{s} = 20$ GeV.



Figure 2: The charm contribution to the proton structure function F_2^c for different values of Q^2 as indicate in the figures. The solid line is the full result, the short-dashed line the hard pomeron contribution and the long-dashed line the soft pomeron contribution. The data are from ZEUS [8] (squares) and H1 [9] (triangles).

It is straightforward to predict the γp total cross section and the $\gamma^* p \to \gamma p$ (DVCS) data, both the Q^2 dependence at fixed $\sqrt{s}$ and the energy dependence at fixed Q^2. All three predictions are in excellent agreement [2] with data. The model predicts a significant contribution from the hard pomeron to the photoproduction cross section, similar to that found in [1]. However the data do not demand a contribution from the hard pomeron due to the comparatively large experimental errors at high energy.

4 $\gamma^{(*)}\gamma^{(*)}$ reactions

With the same approach and the same parameters the $\gamma\gamma$, $\gamma^*\gamma$ (F_2^γ) and $\gamma^*\gamma^*$ cross sections can be obtained. As some of the data are at relatively low energies the Reggeon contribution and the box diagram (the quark-parton-model contribution) have to be included. The Regge contribution was estimated using the form given in [11] and the analytical results of [12] were used for the box diagram. The principal difference between $\gamma\gamma$ and γp at high energies comes



Figure 3: The photon structure function $F_2^\gamma(x,Q^2)/\alpha$ for different values of Q^2. The separate contributions are: soft pomeron (long dashes), hard pomeron (short dashes), box diagram (dot-dashes), Regge (dots). The two solid lines are the combined results including the uncertainty in the Regge term. The data are from OPAL [13] (boxes), L3 [14] (triangles) and ALEPH [15] (stars).

from the singularity of the photon wave function at the origin, which favours the hard-pomeron component. This is apparent even for the scattering of real photons. The energy dependence of the $\gamma\gamma$ total cross section is not compatible with the soft pomeron alone, and the additional contribution of the hard pomeron is required. Once again the predictions of the model are in excellent agreement [2] with the data. The model predictions for F_2^γ/α are equally satisfactory. The comparison with data is made in figure 3. Agreement with experiment is good for small x, but at large x the increasing importance of the Regge term leads to an increasing uncertainty in the predictions. Nonetheless they remain satisfactory for $x < 0.1$. The predictions for $\sigma^{\gamma^*\gamma^*}$ are in fair agreement with the limited data. The only reactions for which the model does not provide a good description of the data are $\gamma^{(*)}\gamma \to c\bar{c}X$, the predictions being about a factor of 2 lower than the L3 [16] cross section for real photons and the OPAL result [17] for the charm contribution to the photon structure function at small x.

5 Conclusions

This simple dipole-dipole model incorporating a soft and a hard pomeron and with only one free parameter provides a remarkably good description of a variety of $\gamma^{(*)}p$ and $\gamma^{(*)}\gamma^{(*)}$ data. The only reactions for which the model does not provide a successful prediction is charm production in $\gamma^{(*)}\gamma$ reactions. In contrast the model does provide an excellent description charm production in $\gamma^{(*)}p$ reactions, explaining the phenomenological result that F_2^c is completely dominated by



Figure 4: The soft-pomeron (solid line) and hard-pomeron (dashed line) contributions to F_2, F_L, F_2^c and F_2^γ/α at different values of x.

the hard pomeron. The relative importance of the soft and hard pomeron contributions to F_2, F_2^c, F_L and F_2^γ are indicated in figure 4. Two features stand out: the dominance of the hard pomeron in F_2^c and the comparability of F_2 and F_2^γ.

A consequence of this approach is that the hard pomeron is not restricted to hard processes, but is present in soft proceses as well. The real $\gamma\gamma$ cross section receives a non-negligible and essential hard contribution due to the point-like coupling of the photon and the resulting singularity of the photon wave function at the origin. There is a hard contribution to the γp cross section, compatible with although not demanded by experiment. There is necessarily a non-zero hard contribution to proton-proton scattering, although at energies presently available it is sufficiently small to be within the limits of experimental error. However at LHC its presence will be observable, although suppressed relative to the simple model by unitarity corrections. Broadly similar conclusions have been obtained in related analyses, although from a somewhat different standpoint, by Cudell *et al* [18] and by Donnachie and Landshoff [19].

References

[1] A. Donnachie and P. V. Landshoff, *Phys. Lett.* B **437**, 408 (1998)

[2] A. Donnachie and H. G. Dosch, *Phys. Rev.* D **65**, 014019 (2002)

[3] O. Nachtmann, Ann. Phys. **209**, 436 (1991)

[4] A. Kraemer and H. G. Dosch, *Phys. Lett.* B **272**, 114 (1991)
H. G. Dosch, E. Ferreira and A. Kraemer, *Phys. Rev.* D **50**, 1992 (1994)

[5] H. G. Dosch, T Gousset, G. Kulzinger and H. J. Pirner, *Phys. Rev.* D **55**, 2602 (1997)
H. G. Dosch, T Gousset and H. J. Pirner, *Phys. Rev.* D **57**, 1666 (1998)

[6] M. Derrick *et al* (ZEUS Collaboration), *Z. Phys.* C **72**, 3 (1996)
J. Breitweg *et al* (ZEUS Collaboration), *Phys. Lett.* B **407**, 432 (1997)

[7] S. Aid *et al* (H1 Collaboration), *Nucl. Phys.* B **470**, 3 (1996)
C. Adloff *et al* (H1 Collaboration), *Nucl. Phys.* B **497**, 432 (1997)

[8] J. Breitweg *et al* (ZEUS Collaboration), *Eur. Phys. J.* C **12**, 35 (2000)

[9] C. Adloff *et al* (H1 Collaboration), *Z. Phys.* C **72**, 593 (1996)

[10] A. Donnachie and P. V. Landshoff, *Phys. Lett.* B **518**, 63 (2001)

[11] A. Donnachie, H. G. Dosch and M. Rueter, *Eur. Phys. J.* C **13**, 141 (2000)

[12] V. M. Budnev *et al*, *Phys.Rep.* **15C**, 182 (1975)

[13] G. Abbiendi *et al* (OPAL Collaboration), *Eur. Phys. J.* C **18**, 15 (2000)
K. Ackerstaff *et al* (OPAL Collaboration), *Phys. Lett.* B **411**, 387 (1997)
K. Ackerstaff *et al* (OPAL Collaboration), *Phys. Lett.* B **412**, 225 (1997)

[14] M. Acciari *et al* (L3 Collaboration), *Phys. Lett.* B **436**, 403 (1997)
M. Acciari *et al* (L3 Collaboration), *Phys. Lett.* B **447**, 147 (1999)

[15] D. Barate *et al* (ALEPH Collaboration), *Phys. Lett.* B **458**, 152 (1999)

[16] M. Acciarri *et al* (L3 Collaboration), *Phys. Lett.* B **453**, 83 (1999); *ibid* **503**, 10 (2001); *ibid*

[17] G. Abbiendi *et al* (OPAL Collaboration), *Eur. Phys. J.* C **16**, 579 (2000)

[18] J. R. Cudell, E. Martynov, O. Selyugin and A. Lengyel, *Phys. Lett.* B **587**, 78 (2004)

[19] A. Donnachie and P.V. Landshoff, *Phys. Lett.* B **595**, 393 (2004)